

# Structure-property modeling of thermodynamic properties of supramolecular systems involving metals or hydrogen bonds

Alexandre Varnek<sup>a</sup>, Vitaly P. Solov'ev<sup>b</sup>

<sup>a</sup> Laboratoire d'Infochimie, Institut de Chimie, 4, rue B. Pascal, Strasbourg 67000, France  
E-mail: varnek@chimie.u-strasbg.fr

<sup>b</sup> Institute of Physical Chemistry, Russian Academy of Sciences, Moscow, Russia  
E-mail: solovev-vp@dio.ru

This presentation concerns the structure – property modeling of the thermodynamic properties of supramolecular systems involving metals or hydrogen bonds. The structure - property studies of existing systems and computer-aided design of new compounds have been performed using the “In Silico” Design and data Analysis (**ISIDA**) system which consists of QSPR, clustering and combinatorial modules as well as some supplementary tools. The **ISIDA** system operates with two types of fragment descriptors: (i) the atom/bond sequences containing from 2 to 15 atoms and (ii) atoms with their nearest environment (“augmented” atoms). Optionally, the fragment descriptors can be mixed with any other “external” descriptors.

The **ISIDA** tools has been successfully tested in modeling of various properties: octanol/water partition coefficients and aqueous solubility of organic molecules, thermodynamic parameters of complexation of metal and neutral guests in solution and in biphasic liquid-liquid systems, biological activities [1-8]. The ensemble of **ISIDA**'s modules has been used for computer-aided design of new binders of some lanthanides and actinides; experimental tests confirms the robustness of predictions. Recently, the assessment of free energy and enthalpy for the 1:1 hydrogen bond complexation of organic acids with organic bases has been performed using an original “labeled atoms” technique. In principle, fragments' contributions to free energy for H-bonding or metal cation – ligand interactions allow one to design new building blocks of supramolecular assemblies possessing desirable binding efficiency.

## References

1. **ISIDA** (In Silico Design and Data Analysis) System.  
<http://infochim.u-strasbg.fr/recherche/isida/index.php>
2. V. P. Solov'ev, A. Varnek, G. Wipff, *Chem. Inf. Comp. Sci.*, 2000, **40** (3), 847-858.
3. A. Varnek, G. Wipff, V. P. Solov'ev *J. Solvent Extract. Ion. Exch.*, 2001, **19** (5), 791 - 837.
4. A. Varnek, G. Wipff, V. P. Solov'ev, A. F. Solotnov *J. Chem. Inf. Comp. Sci.*, 2002, **42** (4), 812-829.
5. V. P. Solov'ev, A. Varnek *J. Chem. Inf. Comp. Sci.*, 2003, **43** (5), 1703 – 1719.
6. A. R. Katritzky, D. C. Fara, H. Yang, M. Karelson, T. Suzuki, V. P. Solov'ev, A. Varnek *J. Chem. Inf. Comp. Sci.*, 2004, **44** (2), 529-541.
7. A. Varnek, D. Fourches, V. P. Solov'ev, V. E. Baulin, A. N. Turanov, V. K. Karandashev, A. R. Katritzky, D. Fara, *J. Chem. Inf. Comp. Sci.*, 2004, **44** (4), 1365 – 1382.
8. V. P. Solov'ev, A. A. Varnek *Rus. Chem. Bul. Int. Ed.* 2004, **53** (7), 1434-1445.